

**ZEEMOX**  
(Amoxicillin)

Capsule: 250mg & 500mg  
Tablet: 1g & 500mg  
Suspension: 125mg/5ml & 250mg/5ml  
Drops: 125mg/1.25ml  
Injection: 250, 500 and 1g

**COMPOSITION**

Zemox 250 & 500 mg Capsule  
Each film coated Capsule contains:  
Amoxicillin trihydrate eq. to Amoxicillin base respectively..... 250 mg & 500 mg (B.P)

Zemox 1g & 500 mg Tablet  
Each film coated Tablet contains:  
Amoxicillin trihydrate eq. to Amoxicillin base respectively..... 1g & 500 mg (U.S.P)

Zemox 125mg/5ml & 250mg/5ml Suspension  
Each 5ml Suspension contains:  
Amoxicillin trihydrate eq. to Amoxicillin base respectively..... 125mg/5ml & 250mg/5ml (B.P)

Zemox 125mg/1.25 ml Drop  
Each 1.25 ml Drop contains:  
Amoxicillin trihydrate eq. to Amoxicillin base respectively..... 125mg/1.25 ml (B.P)

Zemox 250 & 500 mg and 1g injection  
Each vial contains Amoxicillin sodium eq. to Amoxicillin base.....250 mg ,500 mg and 1g (B.P)

**THERAPEUTIC INDICATIONS**

Amoxicillin is indicated for the treatment of the following infections in adults and children.

- Acute bacterial sinusitis
- Acute streptococcal tonsillitis and pharyngitis
- Acute exacerbations of chronic bronchitis
- Community acquired pneumonia
- Acute otitis media
- Acute cystitis
- Acute pyelonephritis
- Asymptomatic Bacteriuria in pregnancy
- Typhoid and paratyphoid fevers
- Dental abscess with spreading cellulitis
- Prosthetic joint infections
- Helicobacter pylori eradication
- Lyme disease

Amoxicillin is also indicated for the prophylaxis of endocarditis. Consideration should be given to official guidance on the appropriate use of antibacterial agents.

**DOSAGE AND ADMINISTRATION**

The dose of Amoxicillin that is selected to treat an individual infection should take into account:

- The expected pathogens and their likely susceptibility to antibacterial agents
- The severity and the site of infection
- The age, weight and renal function of the patient; as shown below

The duration of therapy should be determined by the type of infection and the response of the patient, and should generally be as short as possible. Some infections require longer periods of treatment.

**Adults and children ≥40 kg**

Indication*	Dose*
Acute bacterial sinusitis	250mg to 500mg every 8 hours or 750mg to 1g every 12 hours For severe infections 750mg to 1g every 8 hours Acute cystitis may be treated with 3g twice daily for one day
Asymptomatic bacteriuria in pregnancy	
Acute pyelonephritis	
Dental abscess with spreading cellulitis	
Acute cystitis	500mg every 8 hours, 750mg to 1g every 12 hours For severe infections 750mg to 1g every 8 hours for 10 days
Acute otitis media	
Acute streptococcal tonsillitis and pharyngitis	
Acute exacerbations of chronic bronchitis	
Community acquired pneumonia	500mg to 1g every 8 hours
Typhoid and paratyphoid fever	500mg to 2g every 8 hours
Prosthetic joint infections	500mg to 1g every 8 hours
Prophylaxis of endocarditis	2g orally, single dose 30 to 60 minutes before procedure
Helicobacter pylori eradication	750mg to 1g twice daily in combination with a proton pump inhibitor (e.g. omeprazole, lansoprazole) and another antibiotic (e.g. clarithromycin, metronidazole) for 7 days
Lyme disease	Early stage: 500mg to 1g every 8 hours up to a maximum of 4 g/day in divided doses for 14 days (10 to 21 days) Late stage (systemic involvement): 500mg to 2g every 8 hours up to a maximum of 6 g/day in divided doses for 10 to 30 days
*Consideration should be given to the official treatment guidelines for each indication	

**Children <40 kg**

Children may be treated with Amoxicillin capsules, dispersible tablets, suspensions or sachets.

Amoxicillin Paediatric Suspension is recommended for children under six months of age.

Children weighing 40kg or more should be prescribed the adult dosage.

*Recommended doses:*

Indication*	Dose*
Acute bacterial sinusitis	20 to 90 mg/kg/day in divided doses*
Acute otitis media	
Community acquired pneumonia	
Acute cystitis	
Acute pyelonephritis	
Dental abscess with spreading cellulitis	
Acute streptococcal tonsillitis and pharyngitis	40 to 90 mg/kg/day in divided doses*
Typhoid and paratyphoid fever	100 mg/kg/day in three divided doses
Prophylaxis of endocarditis	50 mg/kg orally, single dose 30 to 60 minutes before procedure
Lyme disease	Early stage: 25 to 50 mg/kg/day in three divided doses for 10 to 21 days Late stage (systemic involvement): 100 mg/kg/day in three divided doses for 10 to 30 days
+ Consideration should be given to the official treatment guidelines for each indication. *Twice daily dosing regimens should only be considered when the dose is in the upper range.	

**Elderly**

No dose adjustment is considered necessary.

### **Renal impairment**

GFR (ml/min)	Adults and children $\geq$ 40kg	Children < 40 kg <sup>#</sup>
greater than 30	No adjustment necessary	No adjustment necessary
10 to 30	Maximum 500mg twice daily	15 mg/kg given twice daily (maximum 500mg twice daily)
less than 10	Maximum 500 mg/day	15 mg/kg given as a single dose (maximum 500 mg)

<sup>#</sup> In the majority of cases, parenteral therapy is preferred

*In patients receiving haemodialysis*

Amoxicillin may be removed from the circulation by haemodialysis

	<b>Haemodialysis</b>
<b>Adults and children over 40kg</b>	500mg every 24h Prior to haemodialysis one additional dose of 500mg should be administered. In order to restore circulating blood levels, another dose of 500mg should be administered after haemodialysis
<b>Children under 40kg</b>	15 mg/kg/day given as a single daily dose (maximum 500mg). Prior to haemodialysis one additional dose of 15 mg/kg should be administered. In order to restore circulating blood levels, another dose of 15 mg/kg should be administered after haemodialysis

*In patients receiving peritoneal dialysis*

Amoxicillin maximum 500mg/day

### **Hepatic impairment**

Dose with caution and monitor hepatic function at regular intervals

#### **Method of Administration**

Amoxicillin is for oral use.

Absorption of amoxicillin is unimpaired by food.

Therapy can be started parenterally according to the dosing recommendations of the intravenous formulation and continued with an oral preparation.

Swallow with water without opening capsule or chewing tablet.

### **CONTRAINDICATIONS**

Hypersensitivity to the active substance, to any of the penicillins or to any of the excipients.

History of a severe immediate hypersensitivity reaction (e.g. anaphylaxis) to another beta-lactam agent (e.g. a cephalosporin, carbapenem or monobactam).

### **SPECIAL WARNINGS AND PRECAUTIONS FOR USE**

#### **Hypersensitivity reactions**

Before initiating therapy with amoxicillin, careful enquiry should be made concerning previous hypersensitivity reactions to penicillin and cephalosporins or other beta-lactam agents.

Serious and occasionally fatal hypersensitivity reactions (including anaphylactoid and severe cutaneous adverse reactions) have been reported in patients on penicillin therapy. These reactions are more likely to occur in individuals with a history of penicillin hypersensitivity and in atopic individuals. If an allergic reaction occurs, amoxicillin therapy must be discontinued and appropriate alternative therapy instituted.

#### **Non-susceptible microorganisms**

Amoxicillin is not suitable for the treatment of some types of infection unless the pathogen is already documented and known to be susceptible or there is a very high likelihood that the pathogen would be suitable for treatment with amoxicillin. This particularly applies when considering the treatment of

patients with urinary tract infections and severe infections of the ear, nose and throat.

#### **Convulsions**

Convulsions may occur in patients with impaired renal function or in those receiving high doses or in patients with predisposing factors (e.g. history of seizures, treated epilepsy or meningeal disorders).

#### **Renal impairment**

In patients with renal impairment, the dose should be adjusted according to the degree of impairment.

#### **Skin reactions**

The occurrence at the treatment initiation of a feverish generalized erythema associated with pustula may be a symptom of acute generalised exanthemous pustulosis. This reaction requires amoxicillin discontinuation and contra-indicates any subsequent administration.

Amoxicillin should be avoided if infectious mononucleosis is suspected since the occurrence of a morbilliform rash has been associated with this condition following the use of amoxicillin.

Patients with lymphatic leukaemia and possibly with HIV infection are particularly prone to developing erythematous rashes with amoxicillin. Amoxicillin should be discontinued if a skin rash occurs.

#### **Jarisch-Herxheimer reaction**

The Jarisch-Herxheimer reaction has been seen following amoxicillin treatment of Lyme disease. It results directly from the bactericidal activity of amoxicillin on the causative bacteria of Lyme disease, the spirochaete *Borrelia burgdorferi*. Patients should be reassured that this is a common and usually self-limiting consequence of antibiotic treatment of Lyme disease.

#### **Overgrowth of non-susceptible microorganisms**

Prolonged use of an anti-infective may result in the overgrowth of non-susceptible organisms (superinfection).

Antibiotic-associated colitis has been reported with nearly all antibacterial agents and may range in severity from mild to life threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhoea during, or subsequent to, the administration of any antibiotics. Should antibiotic-associated colitis occur, amoxicillin should immediately be discontinued, a physician consulted and an appropriate therapy initiated. Anti-peristaltic medicinal products are contra-indicated in this situation.

#### **Prolonged therapy**

Periodic assessment of organ system functions; including renal, hepatic and haematopoietic function is advisable during prolonged therapy. Elevated liver enzymes and changes in blood counts have been reported

#### **Crystalluria**

In patients with reduced urine output, crystalluria has been observed very rarely, predominantly with parenteral therapy. During the administration of high doses of amoxicillin, it is advisable to maintain adequate fluid intake and urinary output in order to reduce the possibility of amoxicillin crystalluria. In patients with bladder catheters, a regular check of patency should be maintained.

#### **Anticoagulants**

Prolongation of prothrombin time has been reported rarely in patients receiving amoxicillin. Appropriate monitoring should be undertaken when anticoagulants are prescribed concomitantly. Adjustments in the dose of oral anticoagulants may be necessary to maintain the desired level of anticoagulation

#### **Interference with diagnostic tests**

Elevated serum and urinary levels of amoxicillin are likely to affect certain laboratory tests. Due to high urinary concentrations of amoxicillin, false positive readings are common with chemical methods.

It is recommended that when testing for the presence of glucose in urine during amoxicillin treatment, enzymatic glucose oxidase methods should be used.

The presence of amoxicillin may distort assay results for oestriol in pregnant women.

### **DRUG INTERACTIONS**

### Oral anticoagulants

Oral anticoagulants and penicillin antibiotics have been widely used in practice without reports of interaction. However, in the literature there are rare cases of increased international normalized ratio in patients maintained on acenocoumarol or warfarin and prescribed a course of amoxicillin. If co-administration is necessary, the prothrombin time or international normalised ratio should be carefully monitored with the addition or withdrawal of amoxicillin. Moreover, adjustments in the dose of oral anticoagulants may be necessary.

### Probenecid

Concomitant use of probenecid is not recommended. Probenecid decreases the renal tubular secretion of amoxicillin. Concurrent use with amoxicillin may result in increased and prolonged blood levels of amoxicillin

### Allopurinol

Concurrent administration of allopurinol during treatment with amoxicillin can increase the likelihood of allergic skin reactions.

### Tetracyclines

Tetracyclines and other bacteriostatic drugs may interfere with the bactericidal effects of amoxicillin.

### Methotrexate

Penicillins may reduce the excretion of methotrexate causing a potential increase in toxicity.

## **FERTILITY, PREGNANCY AND LACTATION**

### Pregnancy:

Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity. Limited data on the use of amoxicillin during pregnancy in humans do not indicate an increased risk of congenital malformations. Amoxicillin may be used in pregnancy when the potential benefits outweigh the potential risks associated with treatment.

### Breastfeeding:

Amoxicillin is excreted into breast milk in small quantities with the possible risk of sensitisation. Consequently, diarrhoea and fungus infection of the mucous membranes are possible in the breast-fed infant, so that breast-feeding might have to be discontinued. Amoxicillin should only be used during breast-feeding after benefit/risk assessment by the physician in charge.

### Fertility:

There are no data on the effects of amoxicillin on fertility in humans. Reproductive studies in animals have shown no effects on fertility.

## **EFFECTS ON ABILITY TO DRIVE AND USE MACHINES**

No studies on the effects on the ability to drive and use machines have been performed. However, undesirable effects may occur (e.g. allergic reactions, dizziness, convulsions), which may influence the ability to drive and use machines

## **ADVERSE DRUG REACTIONS**

The ADRs derived from clinical studies and post-marketing surveillance with amoxicillin, presented by MedDRA System Organ Class are listed below.

The following convention has been utilised for the classification of undesirable effects:

Very common ( $\geq 1/10$ ),

Common ( $\geq 1/100$  to  $< 1/10$ ),

Uncommon ( $\geq 1/1000$  to  $< 1/100$ ),

Rare ( $\geq 1/10,000$  to  $< 1/1000$ ),

Very rare ( $< 1/10,000$ ),

Not known (cannot be estimated from the available data)

The majority of side effects listed below are not unique to amoxicillin and may occur when using other penicillins.

The most commonly reported adverse drug reactions (ADRs) are diarrhoea, nausea and skin rash.

Unless otherwise stated, the frequency of adverse events has been derived from more than 30 years of post-marketing reports.

<b>Infections and infestations</b>	
<b>Very Rare:</b>	Muco-cutaneous candidiasis
<b>Blood and lymphatic system disorders</b>	
<b>Very rare:</b>	Reversible leucopenia (including severe neutropenia or agranulocytosis), reversible thrombocytopenia and haemolytic anaemia. Prolonged prothrombin and bleeding times - Special Warnings and Precautions for Use)
<b>Immune system disorders</b>	
<b>Very rare:</b>	As with other antibiotics, severe allergic reactions, including angioneurotic oedema, anaphylaxis, serum sickness and hypersensitivity vasculitis
<b>Not known:</b>	Jarisch-Herxheimer reaction
If any hypersensitivity reaction occurs the treatment should be discontinued (See also Skin and subcutaneous tissue disorders).	
<b>Nervous system disorders</b>	
<b>Very rare:</b>	Hyperkinesia, dizziness and convulsions. Convulsions may occur in patients with impaired renal function or in those receiving high doses.
<b>Gastrointestinal disorders</b>	
Clinical Trial Data	
<b>*Common:</b>	Diarrhoea and nausea.
<b>*Uncommon:</b>	Vomiting.
Post-marketing Data	
<b>Very rare:</b>	Antibiotic associated colitis (including pseudomembranous colitis and haemorrhagic colitis. Black hairy tongue
<b>Hepato-biliary disorders</b>	
<b>Very rare:</b>	Hepatitis and cholestatic jaundice. A moderate rise in AST and/or ALT.
The significance of a rise in AST and/or ALT is unclear.	
<b>Skin and subcutaneous tissue disorders</b>	
Clinical Trial Data	
<b>*Common:</b>	Skin rash
<b>*Uncommon:</b>	Urticaria and pruritus
Post-marketing Data	
<b>Very rare:</b>	Skin reactions such as erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis, bullous and exfoliative dermatitis, acute generalised exanthematous pustulosis (AGEP) and drug reaction with eosinophilia and systemic symptoms (DRESS).
<b>Renal and urinary tract disorders</b>	
<b>Very rare:</b>	Interstitial nephritis.
<b>Very rare:</b>	Crystalluria.
*The incidence of these AEs was derived from clinical studies involving a total of approximately 6,000 adult and paediatric patients taking amoxicillin.	

## **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at [pv@searlecompany.com](mailto:pv@searlecompany.com)

## **OVERDOSE**

### Symptoms and signs of overdose

Problems of overdosage with amoxicillin are unlikely to occur. Gastrointestinal symptoms (such as nausea, vomiting and diarrhoea) and disturbances of the fluid and electrolyte balances may be evident. Amoxicillin crystalluria, in some cases leading to renal failure, has been observed. Convulsions may occur in patients with impaired renal function or in those receiving high doses.

Treatment of intoxication

Gastrointestinal symptoms may be treated symptomatically, with attention to the water/electrolyte balance.

Amoxicillin can be removed from the circulation by haemodialysis.

## **PHARMACOLOGICAL PROPERTIES**

### **Pharmacodynamic properties**

Pharmacotherapeutic group: penicillins with extended spectrum; ATC code: J01CA04

### Mechanism of action

Amoxicillin is a semisynthetic penicillin (beta-lactam antibiotic) that inhibits one or more enzymes (often referred to as penicillin-binding proteins, PBPs) in the biosynthetic pathway of bactericidal peptidoglycan, which is an integral structural component of the bacterial cell wall. Inhibition of peptidoglycan synthesis leads to weakening of the cell wall, which is usually followed by cell lysis and death.

Amoxicillin is susceptible to degradation by beta-lactamases produced by resistant bacteria and therefore the spectrum of activity of amoxicillin alone does not include organisms which produce these enzymes.

### Pharmacokinetic/ pharmacodynamic relationship

The time above the minimum inhibitory concentration (T>MIC) is considered to be the major determinant of efficacy for amoxicillin.

### Mechanisms of resistance

The main mechanisms of resistance to amoxicillin are:

- Inactivation by bacterial beta-lactamases
- Alteration of PBPs, which reduce the affinity of the antibacterial agent for the target.

Impermeability of bacteria or efflux pump mechanisms may cause or contribute to bacterial resistance, particularly in Gram-negative bacteria.

### Breakpoints

MIC breakpoints for amoxicillin are those of the European Committee on Antimicrobial Susceptibility Testing (EUCAST) version 5.0.

Organism	MIC breakpoint (mg/L)	
	Susceptible ≤	Resistant >
Enterobacteriaceae	8 <sup>1</sup>	8
<i>Staphylococcus</i> spp.	Note <sup>2</sup>	Note <sup>2</sup>
<i>Enterococcus</i> spp. <sup>3</sup>	4	8
Streptococcus groups A, B, C and G	Note <sup>4</sup>	Note <sup>4</sup>
<i>Streptococcus pneumoniae</i>	Note <sup>5</sup>	Note <sup>5</sup>
Viridans group streptococci	0.5	2
<i>Haemophilus influenzae</i>	2 <sup>6</sup>	2 <sup>6</sup>
<i>Moraxella catarrhalis</i>	Note <sup>7</sup>	Note <sup>7</sup>
<i>Neisseria meningitidis</i>	0.125	1
Gram positive anaerobes except <i>Clostridium difficile</i> <sup>8</sup>	4	8
Gram negative anaerobes <sup>8</sup>	0.5	2
<i>Helicobacter pylori</i>	0.125 <sup>9</sup>	0.125 <sup>9</sup>
<i>Pasteurella multocida</i>	1	1
Non-species related breakpoints <sup>10</sup>	2	8

<sup>1</sup>Wild type Enterobacteriaceae are categorised as susceptible to aminopenicillins. Some countries prefer to categorise wild type isolates of *E. coli* and *P. mirabilis* as intermediate. When this is the case, use the MIC breakpoint S ≤ 0.5 mg/L

<sup>2</sup>Most staphylococci are penicillinase producers, which are resistant to amoxicillin. Methicillin resistant isolates are, with few exceptions, resistant to all beta-lactam agents.

<sup>3</sup>Susceptibility to amoxicillin can be inferred from ampicillin

<sup>4</sup>The susceptibility of streptococcus groups A, B, C and G to penicillins is inferred from the benzylpenicillin susceptibility

<sup>5</sup>Breakpoints relate only to non-meningitis isolates. For isolates categorised as intermediate to ampicillin avoid oral treatment with amoxicillin. Susceptibility inferred from the MIC of ampicillin

<sup>6</sup>Breakpoints are based on intravenous administration. Beta-lactamase positive isolates should be reported resistant

<sup>7</sup>Beta lactamase producers should be reported resistant

<sup>8</sup>Susceptibility to amoxicillin can be inferred from benzylpenicillin.

<sup>9</sup>The breakpoints are based on epidemiological cut-off values (ECOFFs), which distinguish wild-type isolates from those with reduced susceptibility.

<sup>10</sup>The non-species related breakpoints are based on doses of at least 0.5g x 3 or 4 doses daily (1.5 to 2 g/day).

The prevalence of resistance may vary geographically and with time for selected species, and local information on resistance is desirable, particularly

when treating severe infections. As necessary, expert advice should be sought when the local prevalence of resistance is such that the utility of the agent in at least some types of infections is questionable.

<b>In vitro susceptibility of micro-organisms to Amoxicillin</b>
<b>Commonly Susceptible Species</b>
<u>Gram-positive aerobes:</u> <i>Enterococcus faecalis</i> Beta-hemolytic streptococci (Groups A, B, C and G ) <i>Listeria monocytogenes</i>
<b><u>Species for which acquired resistance may be a problem</u></b>
<u>Gram-negative aerobes:</u> <i>Escherichia coli</i> <i>Haemophilus influenzae</i> <i>Helicobacter pylori</i> <i>Proteus mirabilis</i> <i>Salmonella typhi</i> <i>Salmonella paratyphi</i> <i>Pasteurella multocida</i>
<u>Gram-positive aerobes:</u> Coagulase negative staphylococcus <i>Staphylococcus aureus</i> <sup>‡</sup> <i>Streptococcus pneumoniae</i> Viridans group streptococcus
<u>Gram-positive anaerobes:</u> <i>Clostridium</i> spp.
<u>Gram-negative anaerobes:</u> <i>Fusobacterium</i> spp.
<u>Other:</u> <i>Borrelia burgdorferi</i>
<b><u>Inherently resistant organisms†</u></b>
<u>Gram-positive aerobes:</u> <i>Enterococcus faecium</i> <sup>†</sup>
<u>Gram-negative aerobes:</u> <i>Acinetobacter</i> spp. <i>Enterobacter</i> spp. <i>Klebsiella</i> spp. <i>Pseudomonas</i> spp.
<u>Gram-negative anaerobes:</u> <i>Bacteroides</i> spp. (many strains of <i>Bacteroides fragilis</i> are resistant)
<u>Others:</u> <i>Chlamydia</i> spp. <i>Mycoplasma</i> spp. <i>Legionella</i> spp.
† Natural intermediate susceptibility in the absence of acquired mechanism of resistance.
‡ Almost all <i>S.aureus</i> are resistant to amoxicillin due to production of penicillinase. In addition, all methicillin-resistant strains are resistant to amoxicillin.

### Pharmacokinetic properties

#### Absorption

Amoxicillin fully dissociates in aqueous solution at physiological pH. Amoxicillin is stable in the acid gastric secretion and is rapidly absorbed from the gastrointestinal tract after oral administration. Following oral administration, amoxicillin is approximately 70% bioavailable. The time to peak plasma concentration (T<sub>max</sub>) is approximately one hour.

The pharmacokinetic results for a study, in which an amoxicillin dose of 250 mg three times daily was administered in the fasting state to groups of healthy volunteers are presented below.

C <sub>max</sub> (µg/ml)	T <sub>max</sub> * (h)	AUC <sub>(0-24h)</sub> (µg.h/ml)	T <sub>½</sub> (h)
3.3 ± 1.12	1.5 (1.0-2.0)	26.7 ± 4.56	1.36 ± 0.56
*Median (range)			

In the range 250 to 3000 the bioavailability is linear in proportion to dose (measured as C<sub>max</sub> and AUC). The absorption is not influenced by simultaneous food intake.

Haemodialysis can be used for elimination of amoxicillin.

#### Distribution

About 18% of total plasma amoxicillin is bound to protein and the apparent volume of distribution is around 0.3 to 0.41/kg. Following intravenous administration, amoxicillin has been found in gall bladder, abdominal tissue, skin, fat, muscle tissue, synovial and peritoneal fluids, bile and pus. Amoxicillin does not adequately distribute into the cerebrospinal fluid.

From animal studies there is no evidence for significant tissue retention of drug-derived material. Amoxicillin, like most penicillins, can be detected in breast milk

Amoxicillin has been shown to cross the placental barrier.

#### Biotransformation

Amoxicillin is partly excreted in the urine as the inactive penicilloic acid in quantities equivalent to up to 10 to 25% of the initial dose.

#### Elimination

The major route of elimination for amoxicillin is via the kidney.

Amoxicillin has a mean elimination half-life of approximately one hour and a mean total clearance of approximately 25 l/hour in healthy subjects. Approximately 60 to 70% of an orally administered dose is excreted unchanged in the urine during the first 6 hours after administration of a single 250mg or 500mg dose of amoxicillin. Various studies have found the urinary excretion to be 50-85% for amoxicillin over a 24-hour period.

Concomitant use of probenecid delays amoxicillin excretion

#### Age

The elimination half-life of amoxicillin is similar for children aged around 3 months to 2 years and older children and adults. For very young children (including preterm newborns) in the first week of life the interval of administration should not exceed twice daily administration due to immaturity of the renal pathway of elimination. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

#### Gender

Following oral administration of amoxicillin to healthy males and female subjects, gender has no significant impact on the pharmacokinetics of amoxicillin.

#### Renal impairment

The total serum clearance of amoxicillin decreases proportionately with decreasing renal function listed

#### Hepatic impairment

Hepatically impaired patients should be dosed with caution and hepatic function monitored at regular intervals.

### **PRECLINICAL SAFETY DATA**

Non-clinical data reveal no special hazard for humans based on studies of safety pharmacology, repeated dose toxicity, genotoxicity and toxicity to reproduction and development.

Carcinogenicity studies have not been conducted with amoxicillin.

### **PRESENTATION**

Zemox 250 Capsule: Pack of 2x10 capsule  
Zemox 500 mg Capsule: Pack of 2x10 capsule  
Zemox 250 mg Capsule: Pack of 10x10 capsule  
Zemox 500 mg Capsule: Pack of 5x10 capsule  
Zemox 500 mg Tablet: Pack of 1x12 tablet  
Zemox 1g mg Tablet: Pack of 2x6 tablet  
Zemox 125mg/5ml Suspension: Pack of 1x90 ml  
Zemox Forte 250mg/5ml Suspension: Pack of 1x90 ml  
Zemox 125mg/1.25 ml Drop: Pack of 1x20 ml  
Zemox injection 1g: Pack of 1 vial  
Zemox injection 250: mg Pack of 1 vial  
Zemox injection 500: mg Pack of 1 vial

### **STORAGE INSTRUCTIONS**

- To be sold on the prescription of a registered medical practitioner only.
- Store in a cool and dry place
- Keep cap tightly closed

- Keep all medicines out of sight & reach of children.

- Shake well before use

- Use within 7 days after reconstitution.

### **REGISTRATION NUMBER**

Zemox 250 Capsule: 015964  
Zemox 500 mg Capsule: 015965  
Zemox 500 mg Tablet: 925472  
Zemox 1g mg Tablet: 017719  
Zemox 125mg/5ml Suspension: 015966  
Zemox 250mg/5ml Suspension: 015967  
Zemox 125mg/1.25 ml Drop: 015968  
Zemox injection 1g:017719  
Zemox injection 250:015970  
Zemox injection 500:015969

### **NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION**

Manufactured by:

ICI Pakistan Ltd.

Pharmaceuticals

32/2A, Phase III, Industrial Estate

Hattar - Pakistan

Marketed by:

The Searle Company Limited,

The Searle Company Limited,

One IBL Centre, 2nd Floor, Plot # 1,

Block 7 & 8, D.M.C.H.S, Tipu Sultan Road,

Off Shakra-e-Faisal, Karachi - Pakistan.

### **DATE OF PUBLICATION OF THE PACKAGE INSERT**

September 2021

SPL/SPC-ZEE.T/C/S/D/921-000(001)